

## REVIEW

Medinformatics

2025, Vol. 00(00) 1–14

DOI: [10.47852/bonviewMEDIN52025024](https://doi.org/10.47852/bonviewMEDIN52025024)



# Biomedicine and Nutritional Epidemiology: AI-Assisted Personalized Nutrition in the Prevention and Management of Psychiatric Disorders

Oluwafikayo Seun Adeyemi-Benson<sup>1,\*</sup>, Rufus Oluwagbemileke Ajayi<sup>1</sup>, Oluwateniola Ajoke Adeyemi-Benson<sup>2</sup> and Taiwo Temitope Ogunjobi<sup>3</sup>

<sup>1</sup>Department of Public Health, University of Illinois at Springfield, USA

<sup>2</sup>Department of Medicine and Surgery, Afe Babalola University, Nigeria

<sup>3</sup>Department of Biochemistry, Ladoké Akintola University of Technology, Nigeria

**Abstract:** Recent evidence emphasizes the critical importance of diet in the prevention and management of psychiatric diseases, including depression, anxiety, schizophrenia, and bipolar disorder. This review examines the significance of personalized nutrition in mental health care within the frameworks of biomedicine and nutritional epidemiology, highlighting its potential to enhance outcomes for individuals with psychiatric disorders. Psychiatric disorders involve complex interactions among genetic, environmental, and lifestyle factors. Although pharmacological treatments are prevalent, personalized nutrition – customizing dietary interventions according to an individual’s genetic composition, microbiome, and metabolic reactions – is gradually recognized as a complementary strategy. Personalized nutrition seeks to enhance cognitive performance and mitigate mental symptoms by using insights from nutrigenomics, epigenetics, and metabolomics. Nutritional epidemiology, which investigates the correlations between diet and health within populations, offers significant insights into dietary patterns and specific nutrients linked to mental health. Epidemiological studies and clinical trials enhance nutritional guidelines for individuals at risk of or managing psychiatric disorders. This review summarizes recent clinical and epidemiological evidence regarding the influence of nutrition on mental health, emphasizing the prospective advantages of personalized nutrition. It also addresses the challenges and future research directions, promoting interdisciplinary methods that integrate biomedicine, nutritional science, and psychiatry to formulate effective, personalized dietary strategies for managing psychiatric disorders.

**Keywords:** personalized nutrition, nutritional epidemiology, psychiatric disorders, biomedicine, mental health, nutrigenomics, mental health management


## 1. Introduction

### 1.1. Overview of psychiatric diseases and their socioeconomic implications

Psychiatric diseases encompass a wide spectrum of mental health conditions characterized by significant changes in an individual’s cognition, emotional regulation, or behavior. Such issues typically lead to substantial distress or impairment in personal, social, and occupational functioning. Among the most prevalent psychiatric disorders are depression, anxiety, bipolar disorder, and schizophrenia, each contributing individually to the global burden of disease. These disorders often have complex causes, including genetic predisposition, environmental influences, and neurobiological abnormalities. Advances in neurology and

psychiatry have enhanced our understanding of the pathogenesis of these illnesses. However, many aspects of their etiology remain unknown, limiting the development of effective treatments. The socioeconomic impact of psychiatric diseases is profound and diverse, affecting individuals, families, and communities (Figure 1) [1]. Mental health disorders are a significant source of impairment worldwide, with depression alone accounting for more than 4% of the global disease burden. The economic repercussions are substantial, with expenditures related to mental health illnesses projected to exceed several trillion dollars annually. These expenses stem from direct healthcare expenditures, loss of productivity, and the economic burden of social assistance programs. Mental problems are frequently linked to heightened morbidity and mortality, in part due to their substantial association with chronic physical conditions like cardiovascular disease, diabetes, and obesity. This bidirectional connection highlights the significance of cohesive strategies in mental and physical health care [2].

\*Corresponding author: Oluwafikayo Seun Adeyemi-Benson, Department of Public Health, University of Illinois at Springfield, USA. Email: [oadey2@uis.edu](mailto:oadey2@uis.edu)



**Figure 1. Global prevalence of major psychiatric disorders (2019)**

In addition to economic and physical concerns, psychiatric disorders profoundly affect psychosocial well-being. Individuals with mental health concerns may encounter stigma and discrimination, exacerbating their struggles and hindering access to help. Notwithstanding heightened public awareness and advocacy initiatives, the societal stigma around psychiatric diseases endures. This stigma not only hinders early identification and treatment but also marginalizes people impacted, limiting their prospects for education, work, and significant relationships. Families and caregivers frequently find themselves entangled in this cycle of pain, enduring emotional burdens, financial pressures, and occasionally social isolation. The repercussions of psychiatric diseases are particularly severe in vulnerable groups, including children, adolescents, and persons living in low- and middle-income countries [3]. The early onset of mental health disorders in teenagers can interrupt their development, leading to poor academic performance, strained relationships, and heightened risks of substance misuse and delinquent conduct. In under-resourced regions, mental health treatments are usually limited, and cultural impediments may further hinder access to care. This gap underlines the global inequalities in mental health resources and emphasizes the importance of international cooperation to solve these concerns [4].

Despite encountering hurdles, psychiatry has made considerable advancements in recent decades. Developments in neuroimaging, genetics, and psychopharmacology have shed light on the molecular roots of mental health concerns, permitting the construction of more tailored and effective therapies. Simultaneously, knowledge of the importance of socioeconomic variables on mental health, including poverty, education, and social support, has expanded. This transition in mental health care supports a more holistic and interdisciplinary approach, incorporating social, psychological, and biological therapies [5]. The impacts of psychiatric diseases transcend beyond individuals who are immediately affected, altering the broader social structure. This issue requires a coordinated effort from governments, healthcare entities, academics, and communities.

Investments in mental health systems, public awareness campaigns, and research are vital for lessening the impact of mental diseases. In addition, developing cultures that boost mental health and eliminate stigma is crucial to building a more inclusive and supportive environment for affected individuals. Mental health should remain a global priority in public health discourse, and measures to address its societal implications must evolve alongside our improving understanding of mental diseases [6]. Figure 1 presents the global prevalence of major psychiatric disorders, including depression, anxiety disorders, bipolar disorder, schizophrenia, and eating disorders, in millions of individuals. The data also includes the percentage of the global population affected by each disorder.

## 1.2. The role of nutrition in mental health

The relationship between nutrition and mental health has become an important research topic, revealing the substantial effect of dietary patterns and individual nutrients on cognitive abilities, emotional stability, and overall mental wellness. Traditionally, mental health research has concentrated primarily on genetic, neurochemical, and psychosocial factors, sometimes missing the essential impact of diet. However, current discoveries underline the relevance of nutrition as a changeable factor in preventing and treating mental diseases such as depression, anxiety, and cognitive loss. This critical shift has led to interdisciplinary research to unravel the biochemical and physiological pathways that relate nutrition to mental health outcomes, thereby opening up new routes for therapy [7]. A central point from this relationship is that the brain requires a continuous supply of nutrients to function efficiently. The vitamins and minerals that support neuroplasticity, neurotransmitter generation, cellular energy activities, and macronutrients like fatty acids are crucial for the brain's structural and functional health. For example, omega-3 fatty acids contained in fish oil are needed for forming brain membranes and have exhibited antidepressant and anti-inflammatory benefits. In addition, reduced levels of B vitamins, particularly

folate, B<sub>6</sub>, and B<sub>12</sub>, are related to changes in homocysteine metabolism, which can interfere with neurotransmitter synthesis and lead to mood problems. Antioxidants, particularly vitamins C and E and polyphenols, help protect against oxidative stress and inflammation, which are increasingly recognized as contributory factors in developing psychiatric illnesses [8].

The emerging area of nutritional psychiatry focuses on how diet affects mental well-being. Research on the Mediterranean diet, typified by a high intake of fruits, vegetables, whole grains, nuts, and olive oil, generally reveals links with lower risks of depression and cognitive impairment. The diet's anti-inflammatory and neuroprotective characteristics are believed to be crucial determinants, confirming the value of a whole-food approach in enhancing mental health. In contrast, Western-style diets, heavy in refined sugars, saturated fats, and processed foods, are associated with lower mental health outcomes, mostly due to their inflammatory effects and bad influence on gut bacteria. The gut-brain axis is an essential part of the nutrition-mental health interaction, a two-way communication pathway connecting the central nervous and gastrointestinal systems. This axis is predominantly governed by gut microbiota, a varied group of bacteria that inhabit the digestive system. Research indicates that dietary choices might impact the makeup and function of gut microbiota, potentially influencing the production of neuroactive chemicals such as gamma-aminobutyric acid (GABA), serotonin, and short-chain fatty acids (SCFAs). Dysbiosis, or an imbalance in gut flora, has been related to mental health conditions such as sadness and anxiety. As a result, treatments that try to restore gut health through prebiotics, probiotics, and fiber-rich diets are gaining attention as prospective strategies to increase mental well-being [10].

The implications of these findings go beyond isolated dietary components, influencing broader public health initiatives and therapeutic practices. Nutritional therapies are increasingly incorporated into mental health care, serving preventive and complementary roles. For example, randomized controlled trials (RCTs) investigating dietary changes alongside standard psychiatric treatments have reported positive outcomes, including mood, cognitive function, and overall quality of life. This evidence reinforces that addressing nutritional deficiencies or imbalances can complement pharmacological and psychotherapy methods, particularly for treatment-resistant individuals [11]. Despite the growing evidence, significant hurdles remain in translating research findings into practice. The heterogeneity in individual dietary reactions shaped by genetic, epigenetic, and metabolic factors highlights the importance of tailored methods in nutritional psychiatry. Advancements in nutrigenomics and metabolomics provide potential ways for personalizing dietary therapy to match individual needs; nevertheless, these technologies are still evolving and need more validation. Socioeconomic problems, such as food insecurity and limited availability to nutrient-rich foods, impede executing dietary guidelines, especially among low- and middle-income populations [12].

### 1.3. Aim of the review

This study studied how a tailored diet is emerging as a possible strategy for treating psychiatric problems. It represents a fascinating juncture where biology, nutritional science, and psychiatry meet in unique ways. The global burden of psychiatric disorders is significant, and there is a pressing need for inventive techniques to enhance treatment outcomes. Personalized nutrition offers a realistic option by adapting dietary interventions to each person's unique genetic makeup, metabolic processes, and gut microbiota

composition. This method examines the biological and environmental variables in psychiatric illnesses. It extensively evaluates the scientific data associating various dietary patterns and individual nutrients with mental health outcomes. By integrating insights from nutrigenomics, epigenetics, and microbiome research, we highlight how individual dietary responses can affect the success of nutritional interventions for psychiatric patients. We also examine the clinical potential of customized dietary plans, drawing on findings from epidemiological studies, clinical trials, and advanced technologies like artificial intelligence (AI)-powered diet optimization and metabolomic profiling. Throughout the review, we critically evaluate the existing research to identify knowledge gaps while discussing the challenges and limitations of implementing personalized nutrition approaches in everyday psychiatric care. By emphasizing the interdisciplinary nature of this work, we foster a deeper understanding of how individualized dietary strategies can be woven into comprehensive mental health treatment plans, ultimately helping to improve outcomes and quality of life for people with psychiatric disorders.

## 2. Understanding the Relationship Between Nutrition and Psychiatric Disorders

### 2.1. Overview of major psychiatric disorders affected by nutrition

Psychiatric disorders are complicated conditions impacted by a range of hereditary, environmental, and lifestyle factors, among which nutrition is becoming acknowledged as a crucial contributor. Emerging research reveals the impact of dietary patterns and specific nutrients on the onset, progression, and severity of various mental diseases, including depression, anxiety, schizophrenia, and bipolar disorder (Table 1). These disorders share standard pathophysiological processes, such as neuroinflammation, oxidative stress, and disruption of the gut-brain axis, all of which are susceptible to dietary treatments. Understanding the role of nutrition in these disorders gives new potential for prevention and management techniques that supplement existing pharmaceutical and psychotherapy approaches [13].

Depression, a prominent cause of disability worldwide, has been extensively investigated in relation to nutrition. Evidence suggests that diets rich in whole foods, such as fruits, vegetables, whole grains, and seafood, are related to a reduced incidence of depression. In contrast, diets high in processed foods, refined carbohydrates, and saturated fats increase vulnerability. The Mediterranean diet, defined by its anti-inflammatory and antioxidant qualities, is particularly beneficial in reducing depression symptoms, perhaps due to its high omega-3 fatty acids, B vitamins, and polyphenols. Omega-3 fatty acids, in particular, perform a key function in modulating neuroinflammation and synaptic plasticity. At the same time, folate and vitamin B<sub>12</sub> shortages impact homocysteine metabolism, decreasing neurotransmitter synthesis and mood modulation [14].

Dietary problems have also been linked to anxiety disorders, including social anxiety disorder, panic disorder, and generalized anxiety disorder. Diet-induced dysregulation of the HPA axis is anticipated to aggravate anxiety symptoms, as stress-related pathways are sensitive to nutritional status. Nutrients such as magnesium, zinc, and tryptophan are essential for maintaining HPA axis homeostasis and promoting the synthesis of GABA, a crucial inhibitory neurotransmitter implicated in anxiety regulation. The gut-brain axis plays a particularly significant role in anxiety disorders, with gut microbiota composition governed by nutrition directly altering brain function through the generation

Table 1. Major psychiatric disorders are influenced by diet, their prevalence, and known dietary factors

Disorder	Global prevalence	Key dietary factors	Impact of diet on disorder
Depression	280 million (3.8%)	Omega-3 fatty acids, B vitamins (B <sub>6</sub> , B <sub>9</sub> , B <sub>12</sub> ), vitamin D, magnesium, antioxidants (e.g., polyphenols)	Nutrient-rich diets reduce inflammation and stimulate neurotransmitter production, relieving depressed symptoms.
Anxiety	301 million (4.0%)	Magnesium, zinc, probiotics (gut-brain axis), omega-3 fatty acids, tryptophan	Supports GABAergic activity and serotonin synthesis, enhancing mood and lowering anxiety symptoms.
Bipolar Disorder	40 million (0.5%)	Omega-3 fatty acids, folate, magnesium, polyunsaturated fatty acids	Stabilizes mood fluctuations and boosts treatment efficacy when paired with pharmaceutical treatments.
Schizophrenia	24 million (0.3%)	Antioxidants (e.g., vitamins C and E), omega-3 fatty acids, glycemic management (low-GI meals)	Reduces oxidative stress and inflammation, which are implicated in the pathogenesis of schizophrenia.
Eating Disorders	14 million (0.2%)	Balanced macronutrient intake, micronutrients (zinc, iron, magnesium), omega-3 fatty acids	Addresses vitamin shortages and helps restore regular eating patterns, boosting overall mental health.

of neuroactive substances and modulation of the immune response. Diets high in fiber and fermented foods, which support healthy gut flora, have been found to lessen anxiety-like behaviors in both clinical and preclinical experiments [15].

Schizophrenia, a severe and chronic psychiatric disorder marked by psychosis, cognitive deficits, and social disengagement, has increasingly been recognized for its ties to dietary variables. Patients with schizophrenia generally tend toward energy-dense, nutrient-poor diets, which contribute to major shortages in key nutrients such as omega-3 fatty acids, antioxidants, and vitamins D and B<sub>12</sub>. These nutritional deficits appear to increase neuroinflammation and oxidative stress, two variables directly associated with the development and progression of schizophrenia. Research suggests that dietary therapies targeting improved glucose management and reduced oxidative damage could ameliorate cognitive deficits and improve overall symptomatology. Auspicious results have emerged from omega-3 fatty acid supplementation, which indicates potential for decreasing the onset of psychosis in high-risk individuals and improving outcomes for those in the early stages of schizophrenia [16].

Bipolar disorder, with its distinctive alternating episodes of mania and sadness, presents particular obstacles in nutritional studies due to its cyclical nature. Nevertheless, accumulating data suggests nutrition plays a considerable role in mood stability and overall disease control. Omega-3 fatty acids have been extensively explored in bipolar disorder, with studies suggesting their potential to lessen depressive episodes and promote mood control. Dysregulated glucose metabolism, a prevalent comorbidity in bipolar disorder, is greatly influenced by dietary choices. High-glycemic-index foods are likely to worsen mood swings and lead to metabolic abnormalities. Furthermore, correlations have been established between abnormalities in magnesium, zinc, and selenium and mood instability, potentially contributing to the underlying pathophysiology of bipolar illness. Findings imply that tailored therapies addressing these nutritional deficits can enhance pharmacological therapy effectiveness and lower the prevalence and severity of mood disorders [17].

The related mechanisms revealed between diet and diverse psychiatric illnesses underline nutrition’s value as a controllable element in mental health treatment. Neuroinflammation, oxidative stress, and gut dysbiosis emerge as significant elements in the pathophysiology of depression, anxiety, schizophrenia, and bipolar disorder, all potentially amenable to dietary management

techniques. While the database grows, greater inquiry is needed to understand the precise biochemical processes through which nutrition affects mental health outcomes. Additionally, tailored techniques incorporating genetic, metabolic, and microbial diversity will be crucial for optimizing nutritional therapy for specific individuals. Incorporating nutrition into comprehensive care plans could potentially benefit quality of life and long-term results for patients affected by these chronic psychiatric diseases [18].

2.2. Mechanisms linking nutrition and mental health

Several biochemical processes, including neuroinflammation, neurotransmitter production, and the gut-brain axis, underpin the complicated link between nutrition and mental health. These pathways describe how dietary components influence brain function and mental well-being, creating a framework for studying the role of nutrition in the origin and progression of psychiatric illnesses. By addressing these pathways, nutritional interventions provide the potential to augment current mental health therapy and give novel strategies for prevention and management [19].

Neuroinflammation constitutes a major mechanism in many psychiatric disorders, showing a vital relationship between food and mental health. The existence of chronic low-grade inflammation in the brain has been related to disorders like depression, anxiety, and cognitive loss. Diet greatly influences this inflammatory response – ingesting pro-inflammatory diets heavy in processed components, trans fats, and refined sweets can aggravate neuroinflammation. On the flip side, anti-inflammatory foods such as those rich in omega-3 fatty acids, polyphenols, and specific vitamins appear to offer protection by reducing microglial activation and decreasing production of pro-inflammatory cytokines like interleukin-6 (IL-6) and tumor necrosis factor-alpha (TNF-α). Omega-3 fatty acids are particularly crucial for resolving inflammation, as they encourage the creation of specialized pro-resolving lipid mediators that help restore balance. Deficiencies in important minerals like magnesium and zinc might increase the inflammatory cascade, underscoring the significance of maintaining a nutrient-dense diet to prevent neuroinflammation [20]. The production of neurotransmitters represents another crucial pathway by which nutrition affects mental health. Neurotransmitters, including serotonin, dopamine, and GABA, play key roles in regulating mood, cognition, and behavior, and



their generation depends on certain dietary precursors and cofactors. Tryptophan, an important amino acid in protein-rich meals, is a precursor for serotonin generation. The brain's access to tryptophan is controlled by food intake and activity in competing metabolic pathways, particularly the kynurenine pathway, which becomes more active during inflammatory conditions. This shift in tryptophan metabolism toward kynurenine lowers serotonin availability, potentially contributing to depressive symptoms. Similarly, tyrosine and phenylalanine, which serve as precursors for dopamine synthesis, depend on adequate food intake and optimal enzyme function, which rely on micronutrients like iron, vitamin B<sub>6</sub>, and vitamin C. The production and release of GABA, the principal inhibitory neurotransmitter, are also impacted by dietary availability, with deficits in magnesium and vitamin B<sub>6</sub> potentially affecting GABAergic function and perhaps leading to anxiety disorders [21].

The gut-brain axis has emerged as a critical mechanism relating nutrition to mental health, demonstrating the two-way communication between the central nervous system and our digestive tract. The gut microbiota, that complex colony of bacteria living in our intestines, plays a critical part in this connection by creating neuroactive chemicals, modulating immunological responses, and changing the integrity of the gut lining. What we consume considerably impacts the composition and variety of these gut microorganisms, with fiber-rich diets boosting the growth of beneficial bacteria that create SCFAs including butyrate, propionate, and acetate. These SCFAs possess anti-inflammatory effects and help maintain the blood-brain barrier, thereby fighting against neuroinflammation [22].

Dysbiosis, an imbalance in gut microbiota, has been linked to psychiatric illnesses such as sadness and anxiety, partly because it increases intestinal permeability and causes systemic inflammation. The gut-brain axis also regulates neurotransmitter production, as specific gut bacteria either directly create serotonin, dopamine, and GABA or change their precursors. For instance, particular strains of *Lactobacillus* and *Bifidobacterium* have been identified to produce GABA, while others modify tryptophan metabolism, influencing serotonin levels. Diets rich in prebiotics and fermented foods can increase the quantity of these beneficial microbes, perhaps providing significant mental health benefits. In contrast, diets high in saturated fats and simple carbohydrates diminish microbial diversity and encourage the proliferation of pro-inflammatory bacteria, contributing to disturbance of the gut-brain axis [23].

Neuroinflammation, neurotransmitter production, and the gut-brain axis don't act in isolation; they're interrelated, generating a complicated web through which nutrition affects mental health. For example, diet-induced abnormalities in gut microbiota can enhance neuroinflammation and impair neurotransmitter synthesis, whereas inflammation can modify gut permeability and microbiota composition. Understanding these associations provides a framework for designing nutritional therapies suited to individual needs, underscoring the need of personalized nutrition methods in mental health care [24].

### 3. Personalized Nutrition

#### 3.1. Scope of personalized nutrition in psychiatric care

Personalized nutrition in psychiatric treatment takes a targeted approach, customizing dietary recommendations and interventions to match each person's unique genetic makeup, metabolic profile, gut microbiome, and environmental factors. This approach recognizes that the connection between what we eat and our mental health varies significantly from person to person,


influenced by genetic variations, epigenetic changes, and the diversity of microbes in our gut. These factors collectively shape how individuals respond to different foods and how these responses affect their mental well-being. By drawing on breakthroughs in fields like nutrigenomics, metabolomics, and microbiome research, personalized nutrition aims to maximize the positive impact of diet on brain function and mental health [12]. The reach of individualized nutrition in mental health therapy encompasses various strategies from identifying specific nutrient deficiencies linked to particular disorders to crafting dietary plans that positively influence neuroinflammation, neurotransmitter production, and gut-brain communication. It also involves using biomarkers to predict how someone might respond to certain foods and integrating digital tools like AI to create adaptive, personalized nutrition programs. This approach holds tremendous promise for addressing the variability seen in psychiatric conditions, where standard treatments often show inconsistent results. Incorporating tailored dietary interventions into psychiatric care offers potential for better symptom control, increased treatment adherence, and stronger long-term mental resilience [25].

#### 3.2. Key components

By integrating individual differences in genetics, epigenetics, microbiome composition, and metabolic profiles, personalized nutrition provides a complete foundation for understanding how dietary treatments can affect mental health. This approach recognizes that our unique physiological and genetic traits influence how we respond to specific nutrients, allowing for individualized dietary regimens that promote mental health advantages. Nutrigenomics studies the relationship between nutrients and an individual's genetic composition, illustrating how genetic variations alter nutritional responses and vulnerability to mental diseases. For example, gene differences in neurotransmitter production or metabolism, such as those affecting folate or serotonin pathways, might drastically modify how specific diets affect mental health. Understanding these genetic distinctions promotes the development of dietary therapies that target specific biological systems, boosting therapeutic outcomes in managing psychiatric diseases [26].

Epigenetics explores how environmental influences, like food, regulate gene expression without modifying the DNA sequence. Nutritional nutrients such as folate, vitamin B<sub>12</sub>, and polyphenols influence epigenetic indicators, including DNA methylation and histone acetylation. These alterations are critical for brain function, neuroplasticity, and resilience to stress. In psychiatric diseases, aberrant epigenetic patterns are widely observed and may be repaired with dietary modifications to restore regular expression and boost mental health results. The microbiome, playing a vital role in the gut-brain axis, regulates the link between diet and mental health. Microbial metabolites, especially SCFAs, impact neuroinflammation, neurotransmitter production, and overall brain function. Variations in microbiome composition, driven by dietary choices, change these pathways and underline the need for individualized microbiome-targeted nutritional treatments. For instance, diets rich in prebiotics and probiotics may boost gut microbial diversity, reduce inflammation, and improve mental health outcomes in disorders like depression and anxiety [27].

Metabolomics complements these components by profiling the biochemical changes occurring after meal consumption. This research helps detect metabolic patterns connected with neurotransmitter pathways, oxidative stress, and inflammation, key components in psychiatric diseases. By evaluating an individual's metabolic profile, tailored nutritional therapy can



**Figure 2. Key aspects of personalized nutrition influencing individual health and well-being**

target particular biochemical abnormalities, enabling more effective management of mental health difficulties. Recent breakthroughs in genetic testing and genomic analysis have considerably advanced the application of tailored nutrition in mental health treatment. These technologies allow the detection of gene variants that alter dietary response, drug metabolism, and illness susceptibility. Such insights give healthcare providers with data to maximize treatment options, including pharmacological regimens and dietary therapy suited to individual needs [28]. Similarly, personalized nutrition leverages analogous technology to build nutritional programs that correspond with an individual's genetic, physiological, and metabolic characteristics, paving the way for precision treatments in controlling psychiatric diseases. Integrating nutrigenomics, epigenetics, microbiome research, and metabolomics creates the cornerstone of customized nutrition. These interwoven parts offer a thorough understanding of the underlying links between nutrition and mental health, creating the foundation for unique, tailored treatments in psychiatric therapy [29]. Figure 2 illustrates the fundamental aspect of customized nutrition, emphasizing that individuals possess distinct physiological and genetic traits that affect their responses to specific nutrients. This information highlights the importance of personalized eating regimens corresponding with these individual biological attributes to maximize mental health outcomes [30].

### 3.3. Genetic makeup and metabolism in dietary responses and mental health

Genetic composition and metabolism together create distinct reactions to diet that greatly affect mental health. The genes that influence nutrient processing, neurotransmitter synthesis, and cellular communication vary widely among persons, explaining the various consequences of dietary alterations and highlighting the need for tailored nutrition methods in mental health therapy. Genetic variants are key to understanding these discrepancies. Specific variants (SNPs) in genes like methylenetetrahydrofolate reductase (MTHFR) and COMT alter critical neurotransmitter pathways. For example, the MTHFR C677T variant impairs folate metabolism, leading to increased homocysteine levels linked to depression and cognitive difficulties. Individuals with this mutation generally respond effectively to specific folate supplementation that helps restore correct methylation and neurotransmitter synthesis. Similarly, changes in genes influencing serotonin synthesis, such as

TPH2, can determine whether eating tryptophan effectively boosts serotonin levels and relieves depression symptoms [26].

Metabolism also determines how nutrition affects mental health by controlling whether nutrients reach the brain and work properly once there. Issues with glucose metabolism are widespread in mental illnesses, with insulin resistance and aberrant brain energy utilization typically reported. Low glycemic index meals can boost mental health by regulating blood sugar and minimizing metabolic stress. Fat metabolism is equally critical in regulating inflammation and cell membrane function, both of which affect neurotransmission. The APOE  $\epsilon 4$  gene variation is associated with altered lipid metabolism and an increased risk for cognitive impairment and mood disorders. Adjusting dietary fat intake to coincide with a person's APOE profile may help attenuate these risks [31].

The relationship between genetics and micronutrient metabolism is critical for mental health. Deficiencies in B vitamins, magnesium, zinc, and iron are common in psychiatric populations and are impacted by hereditary and metabolic variables. Variations in vitamin D receptor genes can impact the effectiveness of vitamin D supplementation for depression. Similarly, changes in iron metabolism genes, such as HFE, can lead to abnormalities in iron levels, which can influence cognitive function and mood regulation. Personalized techniques that combine genetic and metabolic knowledge can optimize micronutrient supplementation for enhanced mental health outcomes [32].

Beyond individual gene effects, polygenic risk scores offer a more comprehensive view by incorporating the influences of several genetic variations associated with psychiatric diseases. These scores can forecast individual dietary demands and weaknesses. Individuals with a high hereditary predisposition for depression might exhibit higher inflammatory reactions to poor diets, making anti-inflammatory foods rich in omega-3s and polyphenols particularly relevant. Metabolomics research exposes distinct metabolic processes connected to psychiatric diseases. Alterations in tryptophan metabolism, with increased activity in the kynurenine pathway, are related to sadness and anxiety. Dietary measures that shift tryptophan toward serotonin synthesis instead of kynurenine may help ease symptoms. Metabolic testing can also indicate abnormalities in fatty acids, amino acids, and antioxidants that can be rectified with focused dietary adjustments [27].

Integrating genetic and metabolic information into nutrition planning represents a transformational approach to mental health care. Advances in nutrigenomics, metabolomics, and personalized medicine allow us to identify individual variances in dietary demands and build specific treatments. By understanding the complicated links between genetic makeup, metabolism, and nutrition, tailored dietary methods may increase the effectiveness of mental health treatment, lessen the burden of illness, and enhance the quality of life for persons with psychiatric illnesses. Further study is needed to confirm these methodologies across varied groups and refine how personalized dietary recommendations are implemented in clinical practice [24].

## 4. Nutritional Epidemiology and Mental Health

### 4.1. Nutritional epidemiology and the diet–mental health relationship

Nutritional epidemiology is a discipline that analyzes the links between dietary patterns, nutrient consumption, and health consequences within populations. It contributes to exploring the connection between nutrition and mental health by identifying linkages, underlying mechanisms, and potential cures. Nutritional epidemiology gives insights into the impact of dietary habits on

the risk and development of psychiatric diseases by evaluating complicated datasets from cohort studies, cross-sectional surveys, and therapy trials. It also plays a significant role in identifying dietary factors that reduce or worsen mental health concerns. Research has frequently linked Mediterranean and anti-inflammatory diets to a decreased risk of depression. Diets heavy in processed foods and refined sugars are related to inferior mental health outcomes [33]. These findings underscore the relevance of dietary quality and specific nutrient elements, such as omega-3 fatty acids, vitamins, and polyphenols, in enhancing mental well-being. Nutritional epidemiology permits examining how individual features, such as genetic predispositions or lifestyle circumstances, affect the relationship between nutrition and mental health. It provides a framework for implementing targeted nutrition therapies and public health policies that increase mental health resilience. By integrating epidemiological data with new disciplines like nutrigenomics and metabolomics, researchers can better grasp the delicate relationship between food and mental health, paving the way for individualized nutrition policies fit for varied populations [8].

## 4.2. Epidemiological evidence connecting diet and psychiatric disorders


Epidemiological research has revealed extensive data tying diet to the occurrence and progression of psychiatric diseases, giving crucial insights into how nutritional patterns influence mental health outcomes. Numerous population-based studies have revealed that those adhering to diets rich in whole foods, such as fruits, vegetables, whole grains, nuts, and fish, tend to have a much-reduced prevalence of sadness, anxiety, and cognitive impairment. These findings underline the preventative impact of nutrient-dense diets like the Mediterranean and DASH, which are rich in anti-inflammatory and neuroprotective compounds. On the other hand, diets heavy in processed foods, trans fats, refined sugars, and low-quality carbs have often been linked to an increased risk of schizophrenia, mood disorders, and even neurodegenerative illnesses. This shows that poor eating habits may increase preexisting susceptibilities to mental health difficulties. Long-term investigations further underscore the causal influence of nutrition on mental health [34]. For example, research

tracking individuals over the years has shown that adherence to Mediterranean or anti-inflammatory diets not only reduces the probability of contracting depressive and anxiety disorders but also mitigates the course of symptoms in those already unwell. This shows that dietary therapies may work as preventive measures and additional treatments to standard psychiatric therapy. Deficiencies in specific nutrients have been linked to increased oxidative stress, neuroinflammation, and impaired neurotransmitter function – processes that are important in many psychiatric diseases. These nutrients include antioxidants like vitamin E and polyphenols, omega-3 fatty acids found in fish, and B vitamins involved in homocysteine regulation (Figure 3) [35].

Epidemiological studies also demonstrate the intricate relationship between diet and other modifiable factors such as lifestyle and socioeconomic conditions. For example, studies have shown that low-income groups experience a disproportionate amount of mental health issues since there are fewer nutrient-dense meals available to them, which exacerbates the psychological effects of food insecurity. Similarly, cultural dietary habits and physical activity levels interact with nutrition to affect mental health outcomes. Traditional diets rich in unprocessed, nutrient-dense foods are commonly connected to superior mental health outcomes, whereas Western dietary patterns are correlated with a heightened risk of psychiatric diseases. Recent research in nutritional epidemiology applies contemporary approaches, including meal pattern analysis and biomarker tests, to find complex correlations. These approaches enable researchers to elucidate the synergistic effects of numerous nutrients and dietary components, rather than concentrated individual nutrients, facilitating a more comprehensive grasp of the diet–mental health relationship [34]. Moreover, integrating genetic and microbiome data in epidemiological research has begun to highlight how individual variances in dietary responses can affect psychiatric outcomes, permitting individualized nutrition in mental health treatment.

### 4.2.1. Omega-3 fatty acids and mental health

Neuronal membranes contain omega-3 fatty acids, particularly docosahexaenoic acid (DHA) and eicosapentaenoic acid (EPA), which are crucial for mental and cognitive health. These important



**Nutrition-related mechanisms in the pathogenesis of mental disorders**

**Figure 3. Nutrition-driven mechanisms in the pathogenesis of mental disorders**



fatty acids retain membrane fluidity, regulate neuroinflammation, and promote neurotransmitter signaling. An increasing amount of evidence indicates that omega-3 fatty acids have preventative effects on depression, anxiety, and other psychiatric diseases. Epidemiological studies suggest that cultures with high dietary intake of omega-3-rich foods, such as fatty fish, display decreased prevalence of depression and mood disorders. Randomized controlled trials have demonstrated the efficacy of omega-3 supplementation, particularly EPA, in lowering depressive symptoms, especially in patients with treatment-resistant depression. The production of pro-inflammatory cytokines, such as IL-6 and TNF- $\alpha$ , which are elevated in depressive illnesses, is known to be suppressed by omega-3 fatty acids. Furthermore, DHA is necessary for synaptic plasticity and cognitive function, suggesting its potential benefits in lowering the risk of neurodegenerative illnesses related to psychiatric disorders [36]. Despite these promising findings, the proper dosage and ratio of EPA to DHA for mental health benefits remain areas of current investigation. Genetic factors that affect the metabolism of omega-3 fatty acids, such as polymorphisms in the fatty acid desaturase gene cluster, may also contribute to individual differences in supplementation responses.

#### 4.2.2. B vitamins and mental health

All these functions, including neurotransmitter synthesis, energy production, and methylation reactions that regulate the brain, require B vitamins, particularly B<sub>6</sub> (pyridoxine), B<sub>9</sub> (folate), and B<sub>12</sub> (cobalamin). A deficiency of these vitamins is frequently linked to depression, cognitive deterioration, and various mental disorders. Folate and B<sub>12</sub> are crucial for maintaining sufficient homocysteine levels, a neurotoxic and inflammatory molecule. Elevated homocysteine levels, typically resulting from insufficient folate or B<sub>12</sub>, are associated with depression and cognitive decline. Supplementation with these vitamins has also been shown to improve depressive symptoms, especially in patients with elevated homocysteine or genetic anomalies influencing folate metabolism, such as the MTHFR C677T polymorphism [36]. Neurotransmitters such as serotonin, dopamine, and GABA are largely produced due to the necessity for vitamin B<sub>6</sub>. Low B<sub>6</sub> levels are linked to decreased neurotransmitter production and, consequently, mood swings and anxiety. B<sub>6</sub> supplementation has also been studied as a potential therapy for several conditions and has proven effective. B vitamins clearly relate to mental health – there's even evidence that, when taken appropriately, they may reduce our risk of dementia and Alzheimer's. However, how well B vitamins promote mental health may depend on baseline nutrition, genetics, and pathology [37].

#### 4.2.3. Antioxidants and mental health

An imbalance between the production of reactive oxygen species and the body's antioxidant defenses is known as oxidative stress, which is increasingly recognized as a contributing factor to mental illnesses. Antioxidants, such as carotenoids, polyphenols, and vitamins C and E, are essential for managing mental health because they prevent oxidative damage and enhance brain function. Vitamin C, a potent antioxidant, is crucial for synthesizing neurotransmitters such as dopamine and norepinephrine. Low levels of vitamin C have been linked to fatigue, depression, and cognitive impairment. Supplementation has been shown to improve mood and reduce anxiety in individuals with insufficient vitamin C levels. The lipid-soluble antioxidant vitamin E also safeguards brain membranes from oxidative damage and is associated with better outcomes in conditions such as neurodegenerative diseases and depression [38]. Polyphenols in foods like berries, tea, and dark chocolate provide neuroprotective benefits by reducing neuroinflammation, enhancing

synaptic plasticity, and modifying signaling pathways involved in mood regulation. Flavonoids, a subclass of polyphenols, have demonstrated promise in enhancing cognitive performance and alleviating symptoms of sadness and anxiety in clinical trials. Carotenoids, particularly lutein and beta-carotene, bolster the brain's antioxidant defenses. Studies have shown a link between higher dietary intake of carotenoids and fewer symptoms of depression and anxiety. Additionally, their anti-inflammatory properties may further benefit mental health by mitigating the impact of chronic systemic inflammation on brain function [39]. Specific nutrients, such as omega-3 fatty acids, B vitamins, and antioxidants, play vital roles in maintaining mental health by reducing neuroinflammation, supporting neurotransmitter function, and protecting against oxidative stress (Table 2). While dietary intake of these nutrients is crucial, supplementation may offer therapeutic benefits, particularly for those with deficiencies or heightened risk factors for psychiatric disorders. Future research should aim to clarify the mechanisms behind these effects and improve nutritional therapy tailored to individual needs, paving the way for personalized nutrition strategies in mental health care [40].

Table 2 gives an overview of significant epidemiological research establishing the relationship between certain nutrients and mental health, stressing the potential for tailored nutritional therapy.

## 5. Personalized Nutritional Interventions

### 5.1. Review of clinical trials and studies on dietary interventions in psychiatric populations

Dietary approaches might also enhance standard treatments for psychiatric disorders, according to clinical trials and studies of diet in psychiatric patients. These treatments often focus on food quality, adding nutrients or modifying dietary patterns, such as the Mediterranean or anti-inflammatory diet. When tailored to patients' needs, these treatments have been shown in RCTs to significantly alleviate symptoms of disorders such as schizophrenia, anxiety, and depression. One area of study is the consumption of omega-3 fatty acids, most commonly EPA, which has been shown to reduce depressive symptoms even in the absence of treatment. Experiments involving other nutrients, including B vitamins, zinc, and magnesium, have suggested varying degrees of effectiveness, generally influenced by baseline deficiencies or genetic factors. The pharmacological effects of folate and B<sub>12</sub> supplements, for instance, have proven effective for individuals with high homocysteine levels.

Whole-diet therapies, such as the Mediterranean diet, have also gained significant interest. These trials often indicate improvements in mood, reductions in anxiety, and enhanced cognitive function, with long-term adherence linked to lasting effects. Mechanistic studies suggest these effects may be mediated by reductions in neuroinflammation, improvements in gut microbiota composition, and enhanced neurogenesis. Despite these promising findings, variation in study designs, sample demographics, and outcomes remains an issue. Future research must focus on refining techniques, integrating biomarkers, and discovering predictors of response to nutritional interventions to develop effective, individualized strategies for addressing psychiatric diseases.

### 5.2. Role of diet in managing specific psychiatric disorders

Diet is essential for managing mental diseases by influencing neuroinflammation, neurotransmitter function, and overall brain health. In cases of depression, adherence to nutrient-dense diets,



Table 2. Summary of epidemiological studies linking specific nutrients to mental health outcomes

Nutrient	Population studied	Key findings	Mental health outcomes
Omega-3 Fatty Acids	Adults with depression	Higher omega-3 consumption associated with lower depressed symptoms; dose-dependent impact seen.	Lower depression severity
B Vitamins (e.g., B <sub>6</sub> , B <sub>12</sub> , Folate)	Elderly individuals	Low levels of B vitamins associated with cognitive impairment and higher risk of depressive disorders.	Enhanced cognitive function; lowered depression risk
Antioxidants (e.g., Vitamin C, E)	General population	High antioxidant consumption correlates with lower stress and anxiety levels.	Lower anxiety and stress symptoms
Zinc	Adolescents with anxiety	Zinc supplementation considerably decreased anxiety symptoms compared to placebo.	Reduce anxiety symptoms
Magnesium	Adults with stress disorders	Low magnesium levels linked to heightened stress and poor mental health effects.	Enhanced stress resilience
Polyphenols (e.g., Flavonoids)	Middle-aged adults	Regular consumption of polyphenol-rich diets associated with enhanced mood and cognitive flexibility.	Improved mood and cognitive performance
Tryptophan	Patients with insomnia	Tryptophan supplementation boosted serotonin synthesis, enhancing sleep and mood.	Improved sleep quality and mood regulation
Vitamin D	Patients with seasonal depression	Vitamin D supplementation lowered symptoms of seasonal affective disorder (SAD).	Lower SAD symptoms

such as the Mediterranean diet, has been shown to reduce symptoms. Essential components like omega-3 fatty acids, B vitamins, and antioxidants aid in mood regulation and lower oxidative stress. Randomized trials have demonstrated that dietary modifications can enhance conventional treatments, particularly in patients with poor baseline diets. For anxiety, the inclusion of foods high in magnesium, zinc, and tryptophan has been associated with reduced symptoms, potentially through their effects on the hypothalamic-pituitary-adrenal axis and serotonin production. Probiotic and prebiotic-rich foods that support gut health have also shown promise in reducing anxiety by altering the gut-brain axis [42]. In cases of schizophrenia, nutritional therapies targeting deficiencies in nutrients such as omega-3 fatty acids and vitamin D have been studied for their ability to improve cognitive performance and alleviate unpleasant symptoms. Although nutritional interventions are not standalone treatments, they offer potential as supplementary measures to enhance overall mental health outcomes in these populations. Emerging research underscores the necessity for personalized nutrition therapies tailored to individual metabolic and genetic profiles for optimal effectiveness [16].

5.3. Personalized nutritional interventions

Personalized nutritional solutions are being thoroughly explored as supplementary techniques for treating psychiatric illnesses, leveraging insights into individual genetic and metabolic variances. This approach acknowledges that responses to food are not uniform but are impacted by genetic variations, metabolic variables, and microbiome composition. Case studies and clinical trials have yielded significant data supporting the utility of personalizing dietary therapy to these qualities, indicating their capacity to benefit mental health outcomes [43].

Case Study 1: MTHFR Gene Polymorphisms and Folate Supplementation in Depression

One of psychiatry’s most well-documented examples of a personalized diet concerns the MTHFR gene, which plays a critical

role in folate metabolism and homocysteine management. Individuals with the MTHFR C677T mutation often display reduced enzyme activity, resulting in increased homocysteine levels and altered methylation pathways critical for neurotransmitter synthesis. Clinical research has shown that patients with this polymorphism enjoy large improvements in depressive symptoms when treated with L-methylfolate, the bioactive form of folate, compared to regular folic acid. For instance, a study conducted on persons with treatment-resistant depression indicated that increased L-methylfolate supplementation boosted response rates and remission, particularly in those with the C677T mutation. This underscores the significance of genotyping in identifying individuals for targeted folate therapy, enabling a personalized strategy to combating mood disorders [44].

Case Study 2: Omega-3 Fatty Acids and APOE Genotypes in Cognitive Decline

The apolipoprotein E (APOE) gene (APOE 4) is a known genetic risk factor for dementia and Alzheimer’s, often connected to neuroinflammation and lipid dysregulation. Omega-3 fatty acid supplementation has also been advised for APOE 4 carriers with poor omega-3 metabolism. In a randomized controlled experiment, APOE 4 carriers fed with high doses of docosahexaenoic acid (DHA) demonstrated reduced cognitive impairment and an enhanced mood compared to those without supplementation. The anti-inflammatory characteristics of omega-3 fatty acids and their ability to modulate neuronal membrane fluidity could potentially lessen the heightened inflammatory response in this gene cluster. This example highlights the significance of genetic testing when developing diets to improve emotional and cognitive well-being [45].

Case Study 3: Gut Microbiota Composition and Probiotic Interventions in Anxiety

The gut-brain axis is also crucial to mental health, and gut microbiota dysbiosis is associated with anxiety and mood disorders. Customized nutritional therapy based on gut microbiota balance, including probiotics and prebiotics, lowered anxiety symptoms. A study of patients with generalized anxiety disorder

indicated a significant reduction in symptoms following a tailored regimen of probiotic strains such as *Lactobacillus rhamnosus* and *Bifidobacterium longum*. The therapy was devised using initial microbiome data that revealed damage to these strains. Gains were related to the synthesis of SCFAs and the regulation of the HPA axis. Personalized probiotic drugs emphasize the potential for microbiome characterization in creating food recommendations for mental health care [46].

#### Case Study 4: Blood Glucose Regulation and Low-Glycemic Diets in Bipolar Disorder

Metabolic dysfunction, such as insulin resistance and poor glucose regulation, is common in bipolar patients and has been linked to mood changes. Individualized diets based on low-glycemic-index (GI) diets have also been tried to regulate mood by modulating blood glucose levels. A controlled trial randomized patients with bipolar disorder and insulin resistance to a low-GI diet that fit their metabolic needs. Such a diet dampened mood swings and increased energy due to the normalization of glucose and insulin. The results highlight the importance of including metabolic variables in individualized nutrition strategies for mental health populations [47].

These case examples illustrate how personalized diet therapies can help navigate the complex interaction between genes, metabolism, and mental health. The treatment is tailored to each patient's genetic variant, metabolism, and microbiome, and personalized nutrition is a practical approach to maximizing

mental health (Table 3). With the development of precision medicine, nutrition in psychiatric care may alter the way mental illness is treated, allowing for more targeted and effective interventions. Future work should aim to develop such methods, utilizing emerging technologies (e.g., genome sequencing and metabolomics) to make them even more precise and effective [48].

Table 3 includes multiple case studies illustrating how personalized dietary regimens, informed by individual genetic and metabolic profiles, can be applied to manage and enhance mental health concerns. These methods involve personalized nutrition and supplements to address specific metabolic pathways implicated in each disease.

#### 5.4. Digital approaches for dietary planning

Advancements in digital health technologies have transformed dietary planning by integrating AI, machine learning, mobile health applications, and personalized nutrition platforms. AI-driven dietary recommendation systems analyze an individual's genomic, metabolic, and behavioral data to create tailored nutritional plans to enhance mental health outcomes. Mobile health (mHealth) applications utilize real-time food intake, physical activity, and psychological well-being tracking to offer adaptive dietary recommendations. Machine learning algorithms are essential in predicting nutrient deficiencies, optimizing meal plans, and recognizing nutritional patterns linked to psychiatric disorders [48]. Wearable devices and smart sensors further improve these

**Table 3. Examples of tailored dietary strategies for depression, anxiety, and schizophrenia based on genetic/metabolic profiles**

Condition	Genetic/Metabolic profile	Tailored dietary strategy	Outcome/effect
Depression	Polymorphisms in MTHFR (methylenetetrahydrofolate reductase) gene influencing folate metabolism	High-dose folate supplementation and B vitamins (B <sub>6</sub> , B <sub>12</sub> ) boost methylation pathways and supports serotonin production.	Significant reduction in depression symptoms, better mood, and cognitive function.
Anxiety	Low serum omega-3 levels and COMT (catechol-O-methyltransferase) gene polymorphism	Omega-3 fatty acid supplementation, particularly EPA (eicosapentaenoic acid), may boost neurotransmitter control and reduce anxiety.	Decreased anxiety symptoms and better stress resilience.
Schizophrenia	Genetic variations in the BDNF (brain-derived neurotrophic factor) gene associated with brain plasticity	Antioxidant-rich diet (e.g., high in flavonoids, vitamin E) to battle oxidative stress and enhance brain function.	Improved cognitive function and fewer psychotic symptoms in schizophrenia.
Depression	Variations in the serotonin transporter (SLC6A4) gene impacting serotonin uptake	Tryptophan-rich diets and supplementation to enhance serotonin levels, along with magnesium supplementation to promote mood control.	Significant mood improvements and fewer depression episodes.
Anxiety	Microbiome dysbiosis and gut-brain axis imbalances	Probiotic and prebiotic supplementation to restore microbial balance, together with a fiber-rich diet, to improve gut health.	Reduced anxiety symptoms and increased gut health, leading to better mental stability.
Schizophrenia	Elevated homocysteine levels due to poor folate metabolism	Folate and B <sub>12</sub> supplements to lower homocysteine levels and support neuronal function, complementing a low-inflammatory diet.	Decreased intensity of schizophrenia symptoms and improved cognitive outcomes.
Depression/Anxiety	Genetic variations in the FKBP5 gene regulating stress response	High-magnesium diet paired with omega-3s to control stress response and lower cortisol levels.	Reduced anxiety and depressed symptoms, increased stress resilience.
Schizophrenia	Genetic mutations in the DRD2 (dopamine receptor D2) gene regulating dopamine signaling	A diet high in zinc and antioxidants to modify dopamine receptor sensitivity and protect against neuroinflammation.	Reduced psychotic symptoms and enhanced cognitive function.

methods by monitoring biomarkers such as glucose levels, heart rate variability, and gut microbiome composition, enabling more data-driven and personalized dietary interventions. Moreover, decision-support tools powered by AI help clinicians and nutritionists design evidence-based, individualized nutritional plans for patients with mental health conditions. Big data analytics integrated with electronic health records facilitate population-wide assessments of dietary habits, contributing to precision nutrition strategies. These digital approaches enhance dietary adherence and engagement and provide scalable, cost-effective solutions for managing psychiatric disorders through personalized nutrition interventions [49].

## 6. Challenges in Implementing Personalized Nutrition

Applying personalized nutrition in clinical practice represents an important advance in health care, delivering more accurate medicines associated with individual genetic, metabolic, and environmental profiles. However, despite its potential, the transition from research to widespread implementation is fraught with challenges and limitations. These include barriers to integration into therapeutic workflows, variability in individual dietary responses, and ethical issues surrounding equitable access and data usage [50].

### 6.1. Barriers to translating personalized nutrition into clinical practice

One of the primary issues in implementing customized nutrition is translating complicated scientific knowledge into practical, scalable, and cost-effective therapies. Current healthcare infrastructures are not designed to accommodate the latest technologies and various information required for customized nutrition. For example, incorporating genetic testing, microbiome analysis, and metabolomics into routine care entails significant expenditures in resources, training, and integration with existing medical systems. Moreover, most healthcare providers lack the professional knowledge to evaluate genetic and metabolic data. This gap underscores the need for collaboration between doctors, dietitians, geneticists, and data scientists. Without such interdisciplinary efforts, the implementation of customized nutrition remains fragmented and inaccessible to the larger community. Financial expense is another key impediment. Investing in sophisticated diagnostics and tailored nutritional therapy typically makes them available only to affluent populations, worsening health disparities. For customized nutrition to become mainstream, cost-reduction methods and policies subsidizing these services are essential [51].

### 6.2. Variability in dietary responses and the need for individualized approaches

Human dietary responses naturally vary and are influenced by genetic polymorphisms, gut microbiota composition, lifestyle, and environmental variables. For example, individuals with specific genetic variations in lipid metabolism-related genes may respond differently to high-fat diets than those without similar variations. Likewise, gut microbial diversity significantly impacts how nutrients are absorbed and utilized. This variation poses a challenge to crafting universally successful personalized feeding regimes. While prediction algorithms utilizing extensive datasets show promise, they often lack the specificity to address individual differences. As a result, interventions that might be effective for one person could have limited or negative effects on another. Furthermore, nutritional therapies require long-term adherence to

achieve meaningful changes. However, adherence can be affected by cultural preferences, economic factors, and personal efficacy assessments. Therefore, personalized nutrition planning must balance scientific accuracy with practical feasibility, employing behavioral strategies to improve compliance [52].

### 6.3. Ethical considerations, including access to personalized care

The ethical concerns of personalized nutrition are enormous, mainly addressing access, privacy, and equity. One key concern is the potential for these measures to aggravate health disparities. Advanced diagnostic techniques and tailored diets are generally prohibitively expensive, limiting their access to affluent individuals. This mismatch risks producing a two-tiered healthcare system, where tailored care becomes a luxury rather than a routine practice. Privacy concerns may arise from collecting and retaining sensitive data, including genomic information and dietary patterns. Ensuring such data's confidentiality and safe treatment is crucial, especially given rising worries about data abuse and breaches [53]. Policies that respect individual rights and specify explicit data-sharing criteria are vital to creating confidence and encouraging participation. Furthermore, ethical difficulties surround the commercialization of personalized nutrition. The increased engagement of private firms in genomic testing and nutritional analytics raises worries about profit-driven ambitions overshadowing scientific rigor and patient welfare. Transparency in the process and evidence underpinning specific concepts is crucial to prevent the growth of pseudoscientific claims. Finally, defining "personalized" treatment poses cultural and ethical difficulties. Dietary advice occasionally overlaps with deeply ingrained cultural behaviors and attitudes. Personalized nutrition techniques must respect and use these cultural traits to avoid alienating specific communities and ensure inclusivity [54].

## 7. Future Directions and Research Opportunities

Future directions in personalized nutrition for mental health necessitate a robust concentration on multidisciplinary research encompassing biomedicine, nutritional science, and psychiatry. This confluence is crucial for understanding the complicated links between nutrition, genetic predispositions, and mental health. For instance, teamwork throughout several professions may show how specific foods influence neurochemical pathways and how dietary responses are shaped by a person's genetic or epigenetic profile. Such integrated research has potential for giving more precise, evidence-based solutions adapted to individual needs [55]. Emerging technologies are set to transform customized nutrition by delivering more detailed evaluations and suggestions. Artificial intelligence, for example, may review enormous volumes of data from genetic, metabolomic, and microbiome research to provide tailored food choices that boost mental health results. Microbiome research is exciting, shining light on the gut-brain axis and its participation in mood regulation. Integrating wearable devices and digital health platforms further augments the capacity to monitor adherence and dynamically adjust dietary recommendations, making interventions more effective and accessible [56]. Promoting public health measures that emphasize the role of nutrition in mental health is also vital. Campaigns to increase awareness about the connection between nutrition and mental well-being could help mitigate stigma and encourage preventive strategies. Meanwhile, policies must focus on providing equitable access to targeted nutrition, especially in disadvantaged regions. Expanding research into individualized mental health treatments

will help bridge the gap between science and practice, enhance clinical outcomes, and foster a more equitable mental health system [57].

The pyramid of evidence levels serves as a fundamental structure in medical research to assess the trustworthiness of scientific results and their therapeutic usefulness. At the pinnacle of the pyramid, systematic reviews and meta-analyses of RCTs supply the most substantial evidence, delivering robust findings based on well-controlled research. Conversely, observational studies, case reports, and expert opinions, which sit in the lowest tiers, are more susceptible to biases and confounding variables, preventing their inclusion into evidence-based therapy guidelines [58]. Most investigations on AI-assisted dietary therapy for psychiatric illnesses often fall within these lower levels of evidence, relying on observational designs, pilot studies, or small-scale trials. While these investigations yield valuable insights, they lack the methodological rigor necessary for strong clinical recommendations [59]. As a result, AI-driven food interventions for mental health remain underrepresented in formal therapy guidelines. To overcome this gap, there is an urgent need for future RCTs and meta-analyses to investigate the efficacy and repeatability of AI-based dietary interventions systematically. By showing causal correlations and collecting statistically accurate data, such research would increase the scientific credibility and therapeutic value of AI-assisted nutritional techniques in psychiatric care. Integrating this hierarchical strategy to evidence review into psychiatric nutrition research exposes the present limitations of dietary interventions in mental health care, stressing the requirement for more rigorous, high-quality investigations. Strengthening the evidence base through well-designed clinical trials will be crucial in encouraging AI-driven dietary interventions as practical components of precision psychiatry and tailored nutrition programs [59].

## 8. Conclusion

In conclusion, the implications of customized nutrition in the context of psychiatric care demonstrate its revolutionary promise in addressing the complex connection between diet and mental disease. This review provides research that relates individual nutrients, eating habits, and specialized interventions to improved outcomes in the treatment of psychiatric diseases, including depression, anxiety, schizophrenia, and bipolar disorder. A molecular perspective on neuroinflammation, neurotransmitter synthesis, and the gut-brain connection provides a biological basis for understanding the significance of customized diets in mental health. Science on nutrigenomics, epigenetics, and the microbiome also emphasizes the necessity of accuracy concerning food and the need for customized interventions. Customized nutrition is a viable strategy to revolutionize psychiatric care by shifting away from generic dietary guidelines toward genetic, metabolic, and lifestyle-dependent approaches. The studies and case reports indicate that treatments targeting specific biomarkers – MTHFR polymorphisms, gut microbiota levels – can enhance mental health outcomes. However, limitations such as food unpredictability, barriers to clinical translation, and questions of access and justice must be overcome to realize its full potential. This highlights a potential role for customized nutrition in psychiatric care, offering a method to bring diet to the forefront of holistic mental health therapy. Making interventions customized, personalized nutrition increases therapeutic effectiveness and empowers patients to engage more actively in their mental health care. As science and technology improve, tailored nutrition in regular psychiatric treatment could transform mental health care by becoming more precise, efficient, and person-centered.

## Acknowledgment

The authors are grateful to the Department of Public Health, University of Illinois at Springfield, USA, for their encouragement and support.

## Ethical Statement

This study does not contain any studies with human or animal subjects performed by any authors.

## Conflicts of Interest

The authors declare that they have no conflicts of interest to this work.

## Data Availability Statement

The data that support this work are available upon reasonable request to the corresponding author.

## Author Contribution Statement

**Oluwafikayo Seun Adeyemi-Benson:** Methodology, Validation, Writing – original draft, Writing – review & editing, Supervision. **Rufus Oluwagbemileke Ajayi:** Conceptualization, Methodology, Software, Investigation. **Oluwatemiola Ajoke Adeyemi-Benson:** Methodology, Visualization, Project administration. **Taiwo Temitope Ogunjobi:** Conceptualization, Methodology, Investigation, Writing – review & editing.

## References

- [1] Forero, D. A. (2023). Genomics of psychiatric disorders: Regional challenges and opportunities. *Biomédica*, 43(1), 5–7. <https://doi.org/10.7705/biomedica.6996>
- [2] Sporinova, B., Manns, B., Tonelli, M., Hemmelgarn, B., MacMaster, F., Mitchell, N., . . . , & Quinn, A. (2019). Association of mental health disorders with health care utilization and costs among adults with chronic disease. *JAMA Network Open*, 2(8), e199910. <https://doi.org/10.1001/jamanetworkopen.2019.9910>
- [3] Ferrie, J., Miller, H., & Hunter, S. C. (2020). Psychosocial outcomes of mental illness stigma in children and adolescents: A mixed-methods systematic review. *Children and Youth Services Review*, 113, 104961. <https://doi.org/10.1016/j.childyouth.2020.104961>
- [4] Rapee, R. M., Oar, E. L., Johnco, C. J., Forbes, M. K., Fardouly, J., Magson, N. R., & Richardson, C. E. (2019). Adolescent development and risk for the onset of social-emotional disorders: A review and conceptual model. *Behaviour Research and Therapy*, 123, 103501. <https://doi.org/10.1016/j.brat.2019.103501>
- [5] Henderson, T. A., Van Lierop, M. J., McLean, M., Uszler, J. M., Thornton, J. F., Siow, Y. H., . . . , & Cohen, P. (2020). Functional neuroimaging in psychiatry—Aiding in diagnosis and guiding treatment. What the American Psychiatric Association does not know. *Frontiers in Psychiatry*, 11, 276. <https://doi.org/10.3389/fpsyt.2020.00276>
- [6] McGorry, P. D., Nelson, B., Wood, S. J., Shah, J. L., Malla, A., & Yung, A. (2020). Transcending false dichotomies and diagnostic silos to reduce disease burden in mental disorders. *Social Psychiatry and Psychiatric Epidemiology*, 55, 1095–1103. <https://doi.org/10.1007/s00127-020-01913-w>



- [7] Horovitz, O. (2025). Nutritional psychology: Review the interplay between nutrition and mental health. *Nutrition Reviews*, 83(3), 562–576. <https://doi.org/10.1093/nutrit/nuae158>
- [8] Muscaritoli, M. (2021). The impact of nutrients on mental health and well-being: Insights from the literature. *Frontiers in Nutrition*, 8, 656290. <https://doi.org/10.3389/fnut.2021.656290>
- [9] Kosyra, K., Drabczyk, M., Marczyńska, Z., Zysk, A., & Magda, I. (2024). Microbiota and depressive disorders—a review. *Journal of Education, Health and Sport*, 60, 188–203. <https://doi.org/10.12775/JEHS.2024.60.013>
- [10] Xiong, R. G., Li, J., Cheng, J., Zhou, D. D., Wu, S. X., Huang, S. Y., . . . , & Li, H. B. (2023). The role of gut microbiota in anxiety, depression, and other mental disorders as well as the protective effects of dietary components. *Nutrients*, 15(14), 3258. <https://doi.org/10.3390/nu15143258>
- [11] Ofor, S. J., Orish, C. N., Frazzoli, C., & Orisakwe, O. E. (2021). Augmenting clinical interventions in psychiatric disorders: Systematic review and update on nutrition. *Frontiers in Psychiatry*, 12, 565583. <https://doi.org/10.3389/fpsyt.2021.565583>
- [12] Wallace, C. (2020). Nutritional psychiatry: A solution for socioeconomic disparities in access to mental health care? *Health Science Inquiry*, 11(1), 155–157. <https://doi.org/10.29173/hsi288>
- [13] Kris-Etherton, P. M., Petersen, K. S., Hibbeln, J. R., Hurley, D., Kolick, V., Peoples, S., . . . , & Woodward-Lopez, G. (2021). Nutrition and behavioral health disorders: Depression and anxiety. *Nutrition Reviews*, 79(3), 247–260. <https://doi.org/10.1093/nutrit/nuaa025>
- [14] Huang, Q., Liu, H., Suzuki, K., Ma, S., & Liu, C. (2019). Linking what we eat to our mood: A review of diet, dietary antioxidants, and depression. *Antioxidants*, 8(9), 376. <https://doi.org/10.3390/antiox8090376>
- [15] Del Toro-Barbosa, M., Hurtado-Romero, A., Garcia-Amezquita, L. E., & García-Cayuela, T. (2020). Psychobiotics: Mechanisms of action, evaluation methods and effectiveness in applications with food products. *Nutrients*, 12(12), 3896. <https://doi.org/10.3390/nu12123896>
- [16] Hassan, N., Dumlao, N., Tran, K., & Zamiri, A. (2022). Improving quality of life with nutritional supplementation in Schizophrenia: A literature review. *European Psychiatry*, 65(S1), S595–S595. <https://doi.org/10.1192/j.eurpsy.2022.1524>
- [17] Choi, J., Kang, J., Kim, T., & Nehs, C. J. (2024). Sleep, mood disorders, and the ketogenic diet: Potential therapeutic targets for bipolar disorder and schizophrenia. *Frontiers in Psychiatry*, 15, 1358578. <https://doi.org/10.3389/fpsyt.2024.1358578>
- [18] Bozzatello, P., Novelli, R., Montemagni, C., Rocca, P., & Bellino, S. (2024). Nutraceuticals in psychiatric disorders: A systematic review. *International Journal of Molecular Sciences*, 25(9), 4824. <https://doi.org/10.3390/ijms25094824>
- [19] Merlo, G., Bachtel, G., & Sugden, S. G. (2024). Gut microbiota, nutrition, and mental health. *Frontiers in Nutrition*, 11, 1337889. <https://doi.org/10.3389/fnut.2024.1337889>
- [20] Businaro, R., Vauzour, D., Sarris, J., Münch, G., Gyengesi, E., Brogelli, L., & Zuzarte, P. (2021). Therapeutic opportunities for food supplements in neurodegenerative disease and depression. *Frontiers in Nutrition*, 8, 669846. <https://doi.org/10.3389/fnut.2021.669846>
- [21] Gasmi, A., Nasreen, A., Menzel, A., Gasmi Benahmed, A., Pivina, L., Noor, S., . . . , & Björklund, G. (2022). Neurotransmitters regulation and food intake: The role of dietary sources in neurotransmission. *Molecules*, 28(1), 210. <https://doi.org/10.3390/molecules28010210>
- [22] Grosso, G. (2021). Nutritional psychiatry: How diet affects brain through gut microbiota. *Nutrients*, 13(4), 1282. <https://doi.org/10.3390/nu13041282>
- [23] Grau-Del Valle, C., Fernández, J., Solá, E., Montoya-Castilla, I., Morillas, C., & Bañuls, C. (2023). Association between gut microbiota and psychiatric disorders: A systematic review. *Frontiers in Psychology*, 14, 1215674. <https://doi.org/10.3389/fpsyg.2023.1215674>
- [24] Estrada, J. A., & Contreras, I. (2019). Nutritional modulation of immune and central nervous system homeostasis: The role of diet in development of neuroinflammation and neurological disease. *Nutrients*, 11(5), 1076. <https://doi.org/10.3390/nu11051076>
- [25] Vieira, C. M., Seidinger Leibovitz, F. M., de Faria Schützer, D. B., Bastos, R. A., Marques de Oliveira, M. R., & Turato, E. R. (2021). A nutritionist approach to psychiatric treatment: Notes from a brief experience. *Revista chilena de nutrición*, 48(5), 798–803. <https://doi.org/10.4067/S0717-75182021000500798>
- [26] Bekdash, R. A. (2024). Epigenetics, nutrition, and the brain: Improving mental health through diet. *International Journal of Molecular Sciences*, 25(7), 4036. <https://doi.org/10.3390/ijms25074036>
- [27] Nohesara, S., Abdolmaleky, H. M., & Thiagalingam, S. (2023). Epigenetic aberrations in major psychiatric diseases related to diet and gut microbiome alterations. *Genes*, 14(7), 1506. <https://doi.org/10.3390/genes14071506>
- [28] Predescu, E., Vaidean, T., Rapciuc, A. M., & Sipos, R. (2024). Metabolomic markers in attention-deficit/hyperactivity disorder (ADHD) among children and adolescents—A systematic review. *International Journal of Molecular Sciences*, 25(8), 4385. <https://doi.org/10.3390/ijms25084385>
- [29] Lagoumintzis, G., & Patrinos, G. P. (2023). Triangulating nutrigenomics, metabolomics and microbiomics toward personalized nutrition and healthy living. *Human Genomics*, 17(1), 109. <https://doi.org/10.1186/s40246-023-00561-w>
- [30] Sevillano-Jiménez, A., Romero-Saldaña, M., García-Rodríguez, M., Molina-Luque, R., & Molina-Recio, G. (2022). Nutritional impact and eating pattern changes in schizophrenic spectrum disorders after health education program on symbiotic dietary modulation offered by specialised psychiatric nursing—two-arm randomised clinical trial. *Nutrients*, 14(24), 5388. <https://doi.org/10.3390/nu14245388>
- [31] Godos, J., Currenti, W., Angelino, D., Mena, P., Castellano, S., Caraci, F., . . . , & Grosso, G. (2020). Diet and mental health: Review of the recent updates on molecular mechanisms. *Antioxidants*, 9(4), 346. <https://doi.org/10.3390/antiox9040346>
- [32] Hui, J., Zhang, N., Kang, M., Gou, Y., Liu, C., Zhou, R., . . . , & Zhang, F. (2024). Micronutrient-associated single nucleotide polymorphism and mental health: A Mendelian randomization study. *Nutrients*, 16(13), 2042. <https://doi.org/10.3390/nu16132042>
- [33] Varela, L. M. (2023). Dietary influence on nutritional epidemiology, public health and our lifestyle. *Nutrients*, 15(11), 2555. <https://doi.org/10.3390/nu15112555>
- [34] Bremner, J. D., Moazzami, K., Wittbrodt, M. T., Nye, J. A., Lima, B. B., Gillespie, C. F., . . . , & Vaccarino, V. (2020). Diet, stress and mental health. *Nutrients*, 12(8), 2428. <https://doi.org/10.3390/nu12082428>
- [35] Dal, N., & Bilici, S. (2024). An overview of the potential role of nutrition in mental disorders in the light of advances in nutripyschiatry. *Current Nutrition Reports*, 13(2), 69–81. <https://doi.org/10.1007/s13668-024-00520-4>
- [36] O'Donovan, F., Carney, S., Kennedy, J., Hayes, H., Pender, N., Boland, F., & Stanton, A. (2019). Associations and effects of

- omega-3 polyunsaturated fatty acids on cognitive function and mood in healthy adults: A protocol for a systematic review of observational and interventional studies. *BMJ open*, 9(6), e027167. <https://doi.org/10.1136/bmjopen-2018-027167>
- [37] Markun, S., Gravestock, I., Jäger, L., Rosemann, T., Pichierri, G., & Burgstaller, J. M. (2021). Effects of vitamin B<sub>12</sub> supplementation on cognitive function, depressive symptoms, and fatigue: A systematic review, meta-analysis, and meta-regression. *Nutrients*, 13(3), 923. <https://doi.org/10.3390/nu13030923>
- [38] Nakao, A., Matsunaga, Y., Hayashida, K., & Takahashi, N. (2021). Role of oxidative stress and Ca<sup>2+</sup> signaling in psychiatric disorders. *Frontiers in Cell and Developmental Biology*, 9, 615569. <https://doi.org/10.3389/fcell.2021.615569>
- [39] Ammar, A., Trabelsi, K., Boukhris, O., Bouaziz, B., Mueller, P., Glenn, J. M., . . . , & Hökelmann, A. (2020). Effects of polyphenol-rich interventions on cognition and brain health in healthy young and middle-aged adults: Systematic review and meta-analysis. *Journal of Clinical Medicine*, 9(5), 1598. <https://doi.org/10.3390/jcm9051598>
- [40] Bozzatello, P., Rocca, P., Mantelli, E., & Bellino, S. (2019). Polyunsaturated fatty acids: What is their role in treatment of psychiatric disorders? *International Journal of Molecular Sciences*, 20(21), 5257. <https://doi.org/10.3390/ijms20215257>
- [41] Dietch, D. M., Kerr-Gaffney, J., Hockey, M., Marx, W., Ruusunen, A., Young, A. H., . . . , & Mondelli, V. (2023). Efficacy of low carbohydrate and ketogenic diets in treating mood and anxiety disorders: Systematic review and implications for clinical practice. *BJPsych Open*, 9(3), e70. <https://doi.org/10.1192/bjo.2023.36>
- [42] Ljungberg, T., Bondza, E., & Lethin, C. (2020). Evidence of the importance of dietary habits regarding depressive symptoms and depression. *International Journal of Environmental Research and Public Health*, 17(5), 1616. <https://doi.org/10.3390/ijerph17051616>
- [43] Wang, F., Zheng, J., Cheng, J., Zou, H., Li, M., Deng, B., . . . , & Kan, J. (2022). Personalized nutrition: A review of genotype-based nutritional supplementation. *Frontiers in Nutrition*, 9, 992986. <https://doi.org/10.3389/fnut.2022.992986>
- [44] Lisboa, J. V. D. C., Ribeiro, M. R., Luna, R. C. P., Lima, R. P. A., do Nascimento, R. A. F., Monteiro, M. G. C. A., . . . , & Costa, M. J. D. C. (2020). Food intervention with folate reduces TNF- $\alpha$  and interleukin levels in overweight and obese women with the MTHFR C677T polymorphism: A randomized trial. *Nutrients*, 12(2), 361. <https://doi.org/10.3390/nu12020361>
- [45] Balakrishnan, J., Husain, M. A., Vachon, A., Chouinard-Watkins, R., Léveillé, P., & Plourde, M. (2022). Omega-3 supplementation increases omega-3 fatty acids in lipid compartments that can be taken up by the brain independent of APOE genotype status: A secondary analysis from a randomised controlled trial 1. *Nutrition and Healthy Aging*, 7(3–4), 147–158. <https://doi.org/10.3233/NHA-220169>
- [46] Yoo, S. Y., Han, A., Park, S., & Lee, J. Y. (2022). Incidence and cognitive decline of Alzheimer's disease and other dementia by apolipoprotein  $\epsilon$ 4 Allele presence: A community-based cohort study in Korean elderly. *Psychiatry Investigation*, 19(3), 190–196. <https://doi.org/10.30773/pi.2021.0347>
- [47] Yassine, H., Arellanes, I. C., Mazmanian, A., De La Cruz, L., Martinez, J., Contreras, L., . . . , & Schneider, L. S. (2023). Baseline findings of PreventE4: A double-blind placebo controlled clinical trial testing high dose DHA in APOE4 carriers before the onset of dementia. *The Journal of Prevention of Alzheimer's Disease*, 10(4), 810–820. <https://doi.org/10.14283/jpad.2023.77>
- [48] Subha, S., Mohankumar, N., Rao, A. S., Khan, A. A., & Murugan, S. (2024). IoT's role in personalized dietary support for mental health using RNN model. In *2024 Third International Conference on Electrical, Electronics, Information and Communication Technologies*, 1–6. <https://doi.org/10.1109/ICEEICT61591.2024.10718428>
- [49] Hickey, B. A., Chalmers, T., Newton, P., Lin, C. T., Sibbritt, D., McLachlan, C. S., . . . , & Lal, S. (2021). Smart devices and wearable technologies to detect and monitor mental health conditions and stress: A systematic review. *Sensors*, 21(10), 3461. <https://doi.org/10.3390/s21103461>
- [50] Pinna, A., & Colasanti, A. (2021). The neurometabolic basis of mood instability: The parvalbumin interneuron link—A systematic review and meta-analysis. *Frontiers in Pharmacology*, 12, 689473. <https://doi.org/10.3389/fphar.2021.689473>
- [51] de Hoogh, I. M., Reinders, M. J., Doets, E. L., Hoevenaars, F. P., & Top, J. L. (2023). Design issues in personalized nutrition advice systems. *Journal of Medical Internet Research*, 25, e37667. <https://doi.org/10.2196/37667>
- [52] Aldubayan, M. A., Pigsborg, K., Gormsen, S. M., Serra, F., Palou, M., Mena, P., . . . , & Hjorth, M. F. (2022). Empowering consumers to PREVENT diet-related diseases through OMICS sciences (PREVENTOMICS): Protocol for a parallel double-blinded randomised intervention trial to investigate biomarker-based nutrition plans for weight loss. *BMJ Open*, 12(3), e051285. <https://doi.org/10.1136/bmjopen-2021-051285>
- [53] Allman-Farinelli, M., Boljevac, B., Vuong, T., & Hekler, E. (2023). Nutrition-related N-of-1 studies warrant further research to provide evidence for dietitians to practice personalized (Precision) medical nutrition therapy: A systematic review. *Nutrients*, 15(7), 1756. <https://doi.org/10.3390/nu15071756>
- [54] Hoevenaars, F. P., Berendsen, C. M., Pasman, W. J., van Den Broek, T. J., Barrat, E., de Hoogh, I. M., & Wopereis, S. (2020). Evaluation of food-intake behavior in a healthy population: Personalized vs. one-size-fits-all. *Nutrients*, 12(9), 2819. <https://doi.org/10.3390/nu12092819>
- [55] Suárez-López, L. M., Bru-Luna, L. M., & Martí-Vilar, M. (2023). Influence of nutrition on mental health: Scoping review. *Healthcare*, 11(15), 2183. <https://doi.org/10.3390/healthcare11152183>
- [56] Park, S. H., Choi, H. K., Park, J. H., & Hwang, J. T. (2024). Current insights into genome-based personalized nutrition technology: A patent review. *Frontiers in Nutrition*, 11, 1346144. <https://doi.org/10.3389/fnut.2024.1346144>
- [57] Grajek, M., Krupa-Kotara, K., Bialek-Dratwa, A., Sobczyk, K., Grot, M., Kowalski, O., & Staśkiewicz, W. (2022). Nutrition and mental health: A review of current knowledge about the impact of diet on mental health. *Frontiers in Nutrition*, 9, 943998. <https://doi.org/10.3389/fnut.2022.943998>
- [58] Banerjee, S., Kumar, D., Kumar, N., Choudhary, N., & Bandyopadhyay, K. (2024). IJCM\_134A: Community engagement strategies for prevention and control of non-communicable diseases in primary healthcare settings: A review of evidences from low-and middle-income countries. *Indian Journal of Community Medicine*, 49, S39. [https://doi.org/10.4103/ijcm.ijcm\\_abstract134](https://doi.org/10.4103/ijcm.ijcm_abstract134)
- [59] Liu, G. D., Li, Y. C., Zhang, W., & Zhang, L. (2020). A brief review of artificial intelligence applications and algorithms for psychiatric disorders. *Engineering*, 6(4), 462–467. <https://doi.org/10.1016/j.eng.2019.06.008>

**How to Cite:** Adeyemi-Benson, O. S., Ajayi, R. O., Adeyemi-Benson, O. A., & Ogunjobi, T. T. (2025). Biomedicine and Nutritional Epidemiology: AI-Assisted Personalized Nutrition in the Prevention and Management of Psychiatric Disorders. *Medinformatics*. <https://doi.org/10.47852/bonviewMEDIN52025024>